DORAMAX INJECTABLE ENDECTOCIDE

ACTIVE CONSTITUENTS: 10 mg/mL Doramectin

SIZE: 500ML

For the treatment and control of doramectin sensitive internal and external parasites of cattle and pigs.

- ✓ Wide safety margin when administered to cattle & pigs.
- ✓ Low irritant formulation, which will minimise any potential reaction to the injection site post administration.
- Broadspectrum treatment for the control of gastrointestinal roundworms, lungworms, sucking lice, biting lice, cattle tick and mange mites.
- The most reliable and accurate dosage format, which ensures animals receive the optimal dose.

DOSE RATE:

CATTLE

1 mL per 50 kg bodyweight, subcutaneously (SC) (under the skin)

1 mL per 33 kg (0.3 mL per 10 kg) bodyweight, intramuscularly (IM)

WITHHOLDING PERIODS:

MEAT

Cattle: DO NOT USE less than 42 days before slaughter for human consumption.

Pigs: DO NOT USE less than 35 days before slaughter for human consumption.

MILK

DO NOT USE in cattle during lactation or less than 60 days before calving when milk or milk products are to be used for human consumption or processing.

STORAGE: Store below 30°C (room temperature).

PACKED IN PLASTIC BOTTLE TO REDUCE CHANCE OF BREAKAGE!

