



DORAMAX

POUR-ON ENDECTOCIDE

BROAD SPECTRUM CONTROL

WIDE SAFETY MARGIN

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

DORAMAX

POUR-ON ENDECTOCIDE
ACTIVE CONSTITUENTS: 5 mg/mL Doramectin

For the treatment and
control of doramectin sensitive
internal and external parasites
of cattle.

1L

2.5L

5L

15L

ACTIVE CONSTITUENT: 5 MG/ML DORAMECTIN

Claims: For the treatment and control of doramectin sensitive internal and external parasites of cattle.

DORAMAX Pour-On Endectocide is for the treatment and persistent control of gastrointestinal roundworms, lungworms and buffalo fly in cattle. Control of eye worms, sucking and biting lice, mange mites and cattle ticks.

DORAMAX Pour-On Endectocide contains doramectin, a member of the macrocyclic lactone family of chemicals.

DORAMAX Pour-On Endectocide is highly effective in the treatment and control of the following species:

Gastrointestinal roundworms (adults and fourth stage larvae)

Adult and immature Lungworm

Sucking Lice

Biting Lice

Buffalo Fly

Cattle Tick

Mange Mites

Dosing and administration:

DOSE RATE: 1 mL of solution per 10 kg bodyweight.

Apply topically along the mid-line of the back in a narrow strip between the withers and tailhead.

WEIGHT (KG)	TREATMENTS PER PACK					
	1L	2.5L	5L	15L	20L	22L
40 – 50	200	500	1000	3000	4000	4400
51 – 75	133	333	666	2000	2666	2933
76 – 100	100	250	500	1500	2000	2200
101 – 150	66	166	333	1000	1333	1466
151 – 200	50	125	250	750	1000	1100
201 – 250	40	100	200	600	800	880
251 – 300	33	83	166	500	666	733
301 – 350	28	71	142	428	571	628
351 – 400	25	62	125	375	500	550
401 – 450	22	55	111	333	444	488
451 – 500	20	50	100	300	400	440
501 – 550	18	45	90	272	360	400
551 – 600	16	41	83	250	333	366
601 – 650	15	38	76	230	307	338

Cattle heavier than 650 kg should be dosed at 1 mL per 10 kg. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on the heaviest cattle in each group (bulls, cows, steers, calves etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight to avoid excessive overdosing.

Treatment of very young animals (less than 40 kg bodyweight) is generally not warranted.

Benefits:

Broad-spectrum: a single treatment with DORAMAX Pour-On Endectocide effectively treats and controls a wide range of internal parasites (roundworms) and external parasites (arthropods) that impair the health and productivity of cattle.

Persistent-action: the persistent drug concentrations that result from the product's unique pharmacokinetics enable it to protect cattle against para- site infection and reinfection for extended periods following treatment.

Safety: DORAMAX Pour-On Endectocide has a wide margin of safety in all classes of cattle. Its broad spectrum of activity, favourable pharmacokinetics and exceptional toleration make it an ideal product for application as an antiparasitic for cattle.

Rainfast: the efficacy of DORAMAX Pour-On Endectocide is not adversely affected if applied when the hide is wet or if rain falls shortly after treatment. However, treatment of animals with wet hides or when rain threatens should be avoided as it is not good agricultural practice.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 42 days before slaughter for human consumption.

MILK: Zero (0) days

TRADE ADVICE – Export Slaughter Interval (ESI):

DO NOT USE less than 42 days before slaughter for export.

For SDS or technical information go to hunterriverco.com.au

APVMA Approval No.: 85729/119780

