

312.5



222

PSU 0972



**BROAD SPECTRUM
ALL WORMER**

10

**CONTROLS
10 PARASITES**



**SINGLE
DOSE**



**EASY
TO ADMINISTER**


CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

ENDURAMEC HORSE PASTE

● **ACTIVE CONSTITUENT: 18.7 g/kg IVERMECTIN**

For treatment and control of roundworms (including arterial larval stages of *Strongylus vulgaris* and benzimidazole-resistant small strongyles), bots, and skin lesions caused by *Habronema* and *Draschia* spp (summer sores) and *Onchocerca* spp microfilariae (cutaneous onchocerciasis).



7.49 g

**ENDURAMEC
HORSE PASTE**

The Hunter River Company Pty Limited | ACN 133 798 615
74-76 Drummond Road, Shepparton VIC 3630 | Ph: (03) 5821 0922



#HUNTER RIVER CO

BATCH:

EXP:



29813C-1

**ENDURAMEC
HORSE PASTE**

ENDURAMEC HORSE PASTE contains ivermectin, a member of the macrocyclic lactones family of chemicals.

INDICATIONS

At the recommended dose level of 200 microgram ivermectin/kg liveweight, ENDURAMEC HORSE PASTE controls the following parasites of horses:

Large Strongyles: *Strongylus vulgaris* (adults and arterial larval stages), *Strongylus edentatus* (adults and tissue stages), *Strongylus equinus* (adults) and *Triodontophorus* spp (adults).

Small Strongyles: Including benzimidazole-resistant strains of adult and immature *Cyathostomum* spp, *Cylicocyclus* spp, *Cylicostephanus* spp, *Cylicodontophorus* spp and *Gyalocephalus* spp.

Pinworm: *Oxyuris equi* (adult and immature)

Ascarids (roundworm): *Parascaris equorum* (adult and immature)

Hairworm: *Trichostrongylus axei* (adult)

Large Mouth Stomach Worm: *Habronema muscae* (adult)

Neck Threadworm: *Onchocerca* spp (microfilariae)

Bots: *Gasterophilus* spp (oral and gastric stages)

Lungworms: *Dictyoacaulus amfieldi* (adult and immature)

Intestinal Threadworm: *Strongyloides westeri* (adult)

ENDURAMEC HORSE PASTE is also effective for control of skin lesions caused by *Habronema* and *Draschia* spp cutaneous larvae (summer sores), and *Onchocerca* spp microfilariae (cutaneous onchocerciasis).

Resistance may develop to any chemical.

DIRECTIONS FOR USE

Contraindications:

This product is contraindicated for use in other animal species as severe adverse reactions may result.

Side Effects:

Some horses have experienced reactions with swelling and itching shortly after treatment. In most cases the horses have been diagnosed as carrying heavy infection of *Onchocerca* microfilariae, and it is assumed the reactions were the result of microfilariae dying in large numbers. Although the signs have resolved in a few days, symptomatic treatment may be advisable. Consult your veterinarian if these clinical signs persist. Resolution of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with ENDURAMEC HORSE PASTE. Reinfection and measures for its prevention should also be considered. Consult your veterinarian if the condition does not improve.

ENDURAMEC HORSE PASTE has been formulated for use in horses only.

Dosage and Administration:

ENDURAMEC HORSE PASTE is given orally at the recommended dose level of 200 µg ivermectin/kg liveweight. The contents of this syringe will treat one horse at 700 kg liveweight. Each weight marking on the syringe plunger will deliver 1.07 g of paste which is sufficient to treat 100 kg liveweight. Remove the plastic cap from the tip of the nozzle. Make sure the horse's

mouth does not contain feed. Insert the syringe into the side of the horse's mouth in the gap between the front and back teeth. Advance the plunger as far as it will go, depositing the medication on the rear of the tongue. Immediately raise the horse's head for a few seconds after dosing to ensure the full dose has been swallowed.

General Directions:

Suggested Parasite Control Program:

All horses should be included in a regular parasite control program with particular attention paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. ENDURAMEC HORSE PASTE is highly effective against gastrointestinal, cutaneous and pulmonary nematodes and bots of horses of all ages. Regular treatment will reduce the chances of verminous arteritis and colic caused by *S. vulgaris*.

With its broad spectrum, ENDURAMEC HORSE PASTE is well suited to be the major product in a parasite control program. Consult your veterinarian for a control program to meet your specific needs.

Product Advantages:

Broad Spectrum: ENDURAMEC HORSE PASTE kills all the important roundworms and arthropod parasites of horses, including the arterial stages of *S. vulgaris* and bots, with a single, low volume, easy to administer dose. Because reinfection is common, treatment must be repeated routinely for effective control.

Safety: ENDURAMEC HORSE PASTE has shown a wide safety margin at the recommended dose level. It may be used in horses of all ages. Mares may be treated at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

MEAT WITHHOLDING PERIODS (HORSES)

DO NOT USE less than 21 days before slaughter for human consumption.
DO NOT USE this preparation in any other animals intended for human consumption.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with the eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ADDITIONAL USER SAFETY INFORMATION

Additional information is listed in the safety data sheet.

ENVIRONMENTAL PROTECTION

Ivermectin is extremely toxic to aquatic species. DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Dispose of empty container by wrapping with paper and putting in garbage.

STORAGE

Store below 30°C (room temperature).

APVMA Approval No. 85529/113512

95

25

255

